

60 YEARS OF NEUROENDOCRINOLOGY

Glucocorticoid dynamics: insights from mathematical, experimental and clinical studies

Francesca Spiga¹, Jamie J Walker^{1,2,3}, Rita Gupta², John R Terry^{2,3} and Stafford L Lightman¹

¹Henry Wellcome Laboratories for Integrative Neuroscience and Endocrinology, University of Bristol, Dorothy Hodgkin Building, Whitson Street, Bristol BS1 3NY, UK

²College of Engineering, Mathematics and Physical Sciences, University of Exeter, Harrison Building, Streatham Campus, North Park Road, Exeter EX4 4QF, UK

³Wellcome Trust Centre for Biomedical Modelling and Analysis, RILD Building, University of Exeter, Exeter, UK

Correspondence should be addressed to S L Lightman
Email
stafford.lightman@bristol.ac.uk

Abstract

A pulsatile pattern of secretion is a characteristic of many hormonal systems, including the glucocorticoid-producing hypothalamic–pituitary–adrenal (HPA) axis. Despite recent evidence supporting its importance for behavioral, neuroendocrine and transcriptional effects of glucocorticoids, there has been a paucity of information regarding the origin of glucocorticoid pulsatility. In this review we discuss the mechanisms regulating pulsatile dynamics of the HPA axis, and how these dynamics become disrupted in disease. Our recent mathematical, experimental and clinical studies show that glucocorticoid pulsatility can be generated and maintained by dynamic processes at the level of the pituitary–adrenal axis, and that an intra-adrenal negative feedback may contribute to these dynamics.

Key Words

- ▶ HPA axis
- ▶ ultradian rhythms
- ▶ glucocorticoids
- ▶ adrenal cortex

Journal of Endocrinology
(2015) 226, T55–T66

Introduction

Glucocorticoids, the end-product of the hypothalamic–pituitary–adrenal (HPA) axis, are essential hormones that regulate the organism's homeostasis and its response to stress. Glucocorticoids (corticosterone in the rat, cortisol in humans, here referred to as CORT) are synthesized in the adrenal cortex in response to adrenocorticotrophic hormone (ACTH) release from corticotroph cells in the anterior pituitary. ACTH secretion is in turn regulated by the release of the neuropeptides corticotropin-releasing hormone (CRH) and arginine vasopressin (AVP) from the paraventricular nucleus of the hypothalamus (PVN). Upon release from the adrenal glands into the general

circulation, CORT exerts its effects by binding specific receptors, the glucocorticoid and mineralocorticoid receptors (GR and MR, respectively), which are widely expressed in target organs throughout the body. In addition to its metabolic, cardiovascular, immune-suppressive and anti-inflammatory effects, CORT also regulates its own production through negative feedback mechanisms within the HPA axis, which include the inhibition of synthesis and release of ACTH from the anterior pituitary (Jones *et al.* 1977), and inhibition of CRH by direct modulation of neuronal activity in the PVN and other brain structures including the hippocampus,

amygdala and prefrontal cortex, which regulate the activity of the PVN (Jones *et al.* 1977, Dallman *et al.* 1987a,b, Ulrich-Lai & Herman 2009) (Fig. 1a).

Ultradian rhythm of the HPA axis

Under basal (i.e., unstressed) conditions, the secretion of CORT over the course of the day is not constant, but is




Figure 1

The HPA axis and glucocorticoid ultradian rhythm. (a) Stress and circadian inputs activate the hypothalamic PVN to release CRH and AVP into the hypothalamic–pituitary portal circulation. CRH and AVP activate corticotroph cells in the anterior pituitary, which respond with the rapid release of ACTH into the general blood circulation. In turn, ACTH reaches the adrenal gland where it activates the synthesis of glucocorticoid hormones (CORT). Glucocorticoids regulate the activity of the HPA axis, and thus their own production, through feedback mechanisms acting at the level of the pituitary gland where they inhibit ACTH release, and at the level of the PVN where they inhibit the release of CRH and AVP. (b) Under basal (i.e., unstressed) conditions, CORT levels are characterized by both a circadian and an ultradian rhythm. The data represent an example 24-h profile of plasma CORT from an adult male rat. Shaded region indicates the dark phase. Adapted from Walker JJ, Spiga F, Waite E, Zhao Z, Kershaw Y, Terry JR & Lightman SL (2012) The origin of glucocorticoid hormone oscillations. *PLoS Biology* 10 e1001341. Published under the terms of the Creative Commons Attribution (CC BY) license.

characterized by a circadian pattern, with hormone levels peaking during the active phase of the animal. In addition to this, studies in several species, including the rat and human, have revealed that CORT is released dynamically from the adrenal gland (Jasper & Engeland 1991, 1994), resulting in an ultradian pulsatile rhythm in the blood (Windle *et al.* 1998), as well as in target tissues, such as the brain (Droste *et al.* 2009), and in subcutaneous tissue (Qian *et al.* 2012). In the rat, CORT pulses have a near-hourly frequency, and changes in the amplitude of these pulses throughout the 24-h cycle determine the circadian variation of hormone secretion (Fig. 1b).

The ultradian rhythm of CORT is an important factor in determining the behavioral, neuroendocrine and genomic response to stressors. Furthermore, CORT pulsatility is crucial for physiological activation of GR and MR and for optimal transcriptional responses of glucocorticoid-responsive genes. Because variation in both the amplitude and frequency of CORT pulses occurs in a number of physiological and pathological conditions, including aging and chronic inflammatory disease (reviewed in Spiga *et al.* (2014)), these changes in the pattern of CORT release may be associated with the disrupted physiological functions observed in these conditions. The importance of pulsatility for genomic, behavioral and neuroendocrine responses to CORT has been described in great detail elsewhere (Spiga *et al.* 2014). The purpose of this review article is to discuss how CORT pulsatility is generated, what the mechanisms are that regulate the dynamics of the HPA axis, and how these dynamics become disrupted in disease.

The origin of glucocorticoid pulsatility

The circadian rhythm of the HPA axis is under the control of the suprachiasmatic nucleus (SCN), which directly regulates the pattern of CRH and AVP release from the PVN, and in addition modulates the responsiveness of the adrenal gland to ACTH via the autonomic nervous system and splanchnic nerve (reviewed in Kalsbeek *et al.* (2012)). There has, however, been much less research into the mechanism underlying the ultradian rhythm of CORT, and how this rhythm is maintained at different levels of the HPA axis, despite the significant amount of data highlighting the importance of the ultradian rhythm. A recent study from our group has demonstrated that while the circadian rhythm of ACTH and CORT is lost in rats in which SCN activity has been disrupted, the ultradian CORT pattern is maintained (Waite *et al.* 2012), providing strong evidence for a CORT ultradian pulse

generator functioning independently of the SCN. This is in accordance with data from Jasper & Engeland (1994) showing that lesioning of the splanchnic nerve results in dampened circadian CORT rhythmicity but sustained pulsatility in adrenal corticosterone secretion measured using intra-adrenal microdialysis techniques.

If the SCN does not generate ultradian rhythmicity, could some other area of the hypothalamus be responsible, as has been described for other pulsatile hormones such as secretion of luteinizing hormone and growth hormone (GH)? These pituitary hormones oscillate in response to the pulsatile secretion of their respective hypothalamic secretagogues (REFS) (Belchetz *et al.* 1978, Clarke & Cummins 1982, Plotsky & Vale 1985).

With respect to HPA axis rhythmicity, pulsatility of ACTH has been described in the rat (Carnes *et al.* 1986, 1988*a,b*), sheep (Apostolakis *et al.* 1992) and human (Henley *et al.* 2009*a*). Furthermore, episodic release of CRH has been shown in macaque (Mershon *et al.* 1992), sheep (Caraty *et al.* 1988, Engler *et al.* 1989), and the rat (Ixart *et al.* 1991, 1994). The evidence of pulsatility of CRH led to the acceptance that the ultradian release of CORT was driven by a hypothalamic CRH pulse generator. There were, however, a number of discrepant findings that challenged this concept. For example, studies in the sheep have shown that ultradian rhythms of ACTH and CORT are maintained even after the hypothalamus has been surgically disconnected from the pituitary (Engler *et al.* 1990), suggesting there must be a hypothalamic-independent pulse generator. Further supporting this idea is the observation that in the rat there is a mismatch between the frequency of CRH pulses (~3 pulses/h) (Ixart *et al.* 1991) and the near-hourly frequency of ACTH and CORT oscillations (Carnes *et al.* 1988*a*).

Using a combination of mathematical modeling and *in vivo* experimental approaches, we have proposed that the pituitary–adrenal system possesses an endogenous oscillatory mechanism generating pulses of ACTH and CORT at a physiological ultradian frequency independent of pulsatile release of CRH and/or AVP from the PVN. This hypothesis is based on the knowledge that the activity of the pituitary–adrenal system is governed by an ACTH-mediated positive feedforward pathway regulating adrenal synthesis and secretion of CORT, and by a CORT-mediated negative feedback that regulates ACTH secretion at the pituitary level. An important feature of this feedforward–feedback loop is the slightly delayed secretory response of the adrenal in response to ACTH stimulation that has been observed both in the rat (Carnes *et al.* 1994, Walker *et al.* 2012) and in humans (Henley *et al.* 2009*a*). This time delay

of a few minutes is presumably due to the time needed by the adrenal for *de novo* synthesis CORT in response to ACTH, since in contrast to peptide hormones (e.g., CRH and ACTH), CORT cannot be pre-synthesized and stored within adrenal steroidogenic cells due to its lipophilic nature. Once released into the general circulation, CORT activates a fast (presumably non-genomic) negative feedback mechanism at the level of the pituitary that still remains to be fully characterized (Jones *et al.* 1972, 1974, Rotsztein *et al.* 1975*a,b*, Mahmoud *et al.* 1984, Widmaier & Dallman 1984, Hinz & Hirschelmann 2000, Russell *et al.* 2010).

Mathematical modeling of the pulse generator

To investigate the dynamic interaction between the pituitary and the adrenal gland, we developed a mathematical model of differential equations incorporating CRH acting on the pituitary, the time taken for CORT to be synthesized and secreted by the adrenal gland in response to ACTH stimulus, and the rapid CORT-mediated inhibition of ACTH secretion at the pituitary (Walker *et al.* 2010). In this study, we made the assumption that CORT-driven inhibition at the level of the anterior pituitary was more important for regulating basal HPA dynamics than CORT feedback at the level of the brain.

We used the mathematical techniques of bifurcation analysis and numerical continuation to study the dynamical behavior of this model (i.e., the dynamics of ACTH and CORT) (Walker *et al.* 2010). These techniques allowed us to identify parameter values – in this case, levels of CRH and adrenal time delay – at which there is a qualitative change in the dynamics of the system (i.e., a bifurcation), and then ‘follow’ this bifurcation through parameter space. This results in curves in parameter space that map out regions corresponding to qualitatively different system dynamics.

In accordance with the hypothesis of a sub-hypothalamic pulse generator, we considered the model’s response to constant levels of CRH drive. The model predicted that for certain ‘physiological’ levels of constant CRH drive (e.g., during the circadian peak of HPA axis activity), the pituitary–adrenal system could support self-sustained ACTH and CORT oscillations characterized by a physiological ultradian frequency (~1 pulse/h; Fig. 2*a*). Furthermore, the model predicted that for either higher ‘stress-equivalent’ levels or lower levels of constant CRH, both ACTH and CORT oscillations become dampened, resulting in a steady-state response in hormone secretion;

such responses have been observed *in vivo* following exposure to an acute stressor (high constant CRH), or at the circadian nadir of HPA axis activity (low constant CRH) (Windle *et al.* 1998).

Interestingly, when a 'physiological level' of pulsatile CRH (~ 3 pulses/h) was imposed on the pituitary–adrenal system, the model predicted that the resulting frequency of ACTH and CORT oscillations (~ 1 pulse/h) would still be predominantly governed by the pituitary–adrenal interaction (Fig. 2b), further supporting an independency of the CORT pulse generator from the dynamics of the hypothalamic drive. Furthermore, when a circadian pattern was imposed on the constant CRH drive, the resultant pattern of ACTH and CORT secretion displayed both circadian and ultradian rhythmicity (Fig. 2c).

To test the validity of our modeling predictions, we then used an experimental approach *in vivo* to investigate the dynamics of ACTH and CORT in response to different levels of constant CRH stimulation in conscious freely behaving male rats (Walker *et al.* 2012). This approach has allowed us to confirm the predictions of our mathematical model, as we have been able to show that ultradian ACTH

and CORT oscillations can be induced by constant infusion of CRH, with resultant CORT pulse amplitude and frequency similar to that of endogenous pulses observed during the circadian peak of hormone release. Consistent with the model, we have also observed the expected time delay between pulses of ACTH and pulses of CORT. Furthermore, constant infusion of a higher dose of CRH resulted in an elevated and sustained level of CORT, similar to the pattern observed in response to severe stressors known to induce a robust increase in hypothalamic CRH secretion.

In summary, both our modeling work and experimental data support the hypothesis of a self-sustained, sub-hypothalamic ACTH and CORT pulse generator that depends on the dynamic interaction between the anterior pituitary and the adrenal cortex.

Our mathematical model also predicts that any disruption in the ACTH feedforward drive (i.e., adrenal delay) or in the CORT feedback mechanism within the pituitary could have an effect on the pulsatile dynamics of ACTH and CORT. Since ACTH is not the only factor that can influence CORT synthesis in the adrenal, it is clearly important to investigate how other factors such as pathogens and inflammatory mediators may interact with the physiological ACTH-mediated adrenal response. Chronic inflammation in the rat is associated with changes in the pulsatile CORT pattern (Windle *et al.* 2001). Furthermore, changes in CORT-mediated signaling at the level of the anterior pituitary can also affect CORT pulsatility, and chronic administration of a GR antagonist leads to changes in the ultradian rhythm of CORT with an increase in the number, height and frequency of CORT pulses, resulting in an overall increase in hormone levels through the 24-h cycle, presumably as a result of an impaired GR-mediated inhibition of ACTH secretion (Spiga *et al.* 2007).




Figure 2

CORT responses (black) to different patterns of CRH drive (grey). Model predictions of CORT pulsatility for constant CRH (a), pulsatile CRH (b) and circadian CRH (c). Response demonstrates a frequency in CORT governed by the pituitary–adrenal system and not by the frequency of the CRH forcing. Data adapted from Walker *et al.* (2010). Adapted from Walker JJ, Terry JR & Lightman SL (2010) Origin of ultradian pulsatility in the hypothalamic–pituitary–adrenal axis. *Proceedings of the Royal Society B: Biological Sciences* 277 1627–1633. Published under the terms of the Creative Commons Attribution (CC BY) license.

Adrenal regulation of glucocorticoid pulsatility

Studies in human and in the rat have shown that in addition to CORT, ACTH is also released in a pulsatile manner (Fig. 3D), with the ACTH pulse amplitude increasing throughout the day as result of the circadian input from the SCN, via regulation of circadian secretion of CRH (Watts *et al.* 2004). Given that pulsatility of CORT is important for maintaining optimal transcription of glucocorticoid regulated genes (Stavreva *et al.* 2009), we decided to investigate whether pulsatile ACTH is important for optimal transcription of ACTH-responsive




Figure 3

Dynamics of human ACTH and cortisol in health and disease. Changes in cortisol and adrenocorticotropic hormone (ACTH) levels throughout the 24-h perioperative period of cardiac surgery (A–C), and in a healthy volunteer (D). (A) mean \pm s.e.m. 24-h cortisol and ACTH profile from patients undergoing coronary artery bypass grafting (CABG). (B) specific mean \pm s.e.m. 24-h cortisol profile from patients undergoing CABG using the off-pump or the on-pump technique. In (A) and (B) the light grey area represents the period during which some patients were undergoing surgery. The dark grey area represents the period during which all patients were undergoing surgery. (C) individual 24-h ACTH and cortisol profile of a patient undergoing off-pump CABG; the light grey area represents the period during which the patient was undergoing surgery (09:19–13:49 h). After the initial surge of ACTH and cortisol, both ACTH and cortisol continue to pulse. However, while both the absolute values of ACTH and the pulse amplitude are reduced, the cortisol levels remain elevated. The grey area represents the period of surgery. (D) individual 24-h ACTH and cortisol profile from a healthy volunteer. ACTH and cortisol both display a tightly correlated ultradian rhythm. Reproduced, with permission, from Gibbison B, Spiga F, Walker JJ, Russell GM, Stevenson K, Kershaw Y, Zhao Z, Henley D, Angelini GD & Lightman SL (2015) Dynamic pituitary-adrenal interactions in response to cardiac surgery. *Critical Care Medicine* 43 791–800. Copyright 2015 Society of Critical Care Medicine and Wolters Kluwer Health, Inc. The original source of (D) is Henley DE, Leendertz JA, Russell GM, Wood SA, Taheri S, Woltersdorf WW & Lightman SL, *Journal of Medical Engineering & Technology*, 2009; 33: 199–208, copyright 2009 Informa Healthcare. Reproduced with permission of Informa Healthcare.

steroidogenic genes within the adrenal cortex of the rat. By using a model of suppressed endogenous ACTH secretion, we found that while the adrenal gland responds rapidly to pulses of ACTH, with pulsatile activation of the steroidogenic pathway *in vivo* that parallels pulsatile secretion of CORT, this dynamic response is absent when an identical dose of ACTH is infused at a constant rate (Spiga *et al.* 2011a). Moreover, the responsiveness of the adrenal gland to a pulse of ACTH equivalent to that seen during an acute stress response is also reduced in rats infused with constant ACTH (Spiga & Lightman 2015). These observations suggest that the adrenal is adapted to respond rapidly to individual pulses of ACTH, and that optimal adrenal responsiveness depends on pulsatile ACTH. We have now begun to dissect the mechanisms underlying this phenomenon by investigation of the dynamics of the adrenal steroidogenic pathway.

ACTH induces CORT synthesis by activating its specific cell surface G-protein-coupled receptor, the melanocortin type-2 receptor (MC2R; Mountjoy *et al.* 1994). ACTH binding to MC2R leads to activation of adenylyl cyclase, followed by an increase in intracellular levels of cAMP, which in turn activates downstream signaling pathways including the protein kinase A (PKA) pathway. Since CORT cannot be stored within adrenal cells, the rapid synthesis of CORT that occurs within minutes upon ACTH stimulation *in vivo* must depend on ACTH-induced non-genomic events that include post-translational modification of steroidogenic proteins. This includes phosphorylation of proteins involved in cholesterol metabolism, notably including hormone-sensitive lipase (HSL) and steroidogenic acute regulatory protein (StAR), which regulate the levels of intracellular cholesterol and its transport within the mitochondria matrix,

respectively (Lin *et al.* 1995, Kraemer & Shen 2002). In addition to these rapid non-genomic events, ACTH also regulates the transcription of genes encoding for steroidogenic proteins, including *StAR*, *CYP11A* (the gene encoding for the cholesterol side-chain cleavage cytochrome protein P450_{scc} (Churchill & Kimura 1979), which catalyses the cleavage of the cholesterol side-chain to produce pregnenolone in the mitochondria) and *MRAP* (the gene encoding for the melanocortin receptor accessory protein (MRAP), which regulates the level and activity of MC2R at the cell surface, and thus the cell's responsiveness to ACTH; Metherell *et al.* 2005). We have shown that *StAR*, *CYP11A* and *MRAP* mRNA are normal in rats infused with pulsatile ACTH, but actually decrease in rats infused with constant ACTH (Spiga *et al.* 2011a).

Studies investigating the effects of a single ultradian pulse of ACTH on the dynamics of steroidogenic gene transcription in the rat show that the adrenal steroidogenic pathway is highly dynamic in response to a pulse of ACTH, with rapid changes in the levels of *StAR*, *CYP11A1* and *MRAP* transcription, measured as changes in levels of heteronuclear RNA (hnRNA), peaking 15 min after ACTH administration and returning to basal levels within 30 min (Spiga *et al.* 2011b, Liu *et al.* 2013). It is well known that transcription of *StAR* and other steroidogenic genes is regulated by the phosphorylation of CREB and subsequent CREB binding to the CRE with the target gene promoter. Consistent with this, we found that the dynamic transcriptional activation of *StAR* in response to a pulse of ACTH was associated with rapid and transient phosphorylation of CREB and rapid dephosphorylation and nuclear translocation of the CREB co-activator, transducer of regulated CREB activity 2 (TORC2, also called CRTC2) (Takemori *et al.* 2007), both occurring 5 min after administration of the ACTH pulse.

These findings show that pulsatile events leading to the dynamic transcription of steroidogenic genes in the adrenal gland parallel the pulsatile secretion of CORT. We propose that pulsatile ACTH results in pulsatile expression of *StAR*, and other steroidogenic genes such as *MRAP*, throughout the circadian cycle, and therefore increases in both the amplitude of ACTH pulses and/or responsiveness of the adrenal to ACTH during the circadian peak will lead to increased amplitude in pulsatile transcription of steroidogenic genes, ultimately resulting in the circadian variation in steroidogenic protein levels that we have observed (Fig. 4) (Park *et al.* 2013). In addition to *StAR*, several transcription factors involved in *StAR*'s transcriptional regulation, including the positive regulators steroidogenic factor 1 (SF-1; Caron *et al.* 1997) and Nur77




Figure 4

Schematic model of dynamic expression of steroidogenic genes. Pulsatile ACTH induces dynamic transcription of *StAR* (hnRNA); circadian variation in ACTH pulse amplitude, as well as circadian variation in adrenal responsiveness to ACTH, determine changes in the amplitude of *StAR* hnRNA pulses leading to circadian expression of *StAR* protein. Model based on data from Carnes *et al.* (1988a), Spiga *et al.* (2011b) and Park *et al.* (2013).

(NR4A1; Martin *et al.* 2008) and the negative regulator DAX-1 (dosage sensitive sex-reversal (DSS), adrenal hypoplasia congenita (AHC) locus on the X-chromosome; Jo & Stocco 2004), are also rhythmically expressed in the adrenal gland (Park *et al.* 2013). The pattern of expression of SF-1, Nur77 and DAX-1 is consistent with their role in regulating the expression of adrenal *StAR*: while SF-1 and Nur77 mRNA and protein levels are elevated when *StAR* expression reaches its circadian peak, the levels of expression of DAX-1 are high when *StAR* expression is low. Given that ACTH inhibits DAX-1 transcription, the circadian increase in ACTH levels decreases DAX-1 expression, thus reducing its inhibitory effect on *StAR* transcription. The resulting increase in *StAR* expression in turn contributes to increased CORT levels during the circadian peak. It is interesting that in rats in which the CORT circadian rhythm has been disrupted by chronic exposure to constant light, the circadian rhythm of *StAR* and its transcriptional regulators is also disrupted (Park *et al.* 2013), further supporting our hypothesis that the pattern of steroidogenic activity is crucial for maintaining optimal rhythmicity of CORT secretion.

Intra-adrenal glucocorticoid-mediated negative feedback

Our work implicates a key role for rapid CORT feedback of CRH-induced ACTH secretion from the pituitary in regulating the ultradian activity of the system. Since the adrenal cortex expresses GR, as shown by studies both in rodents (Loose *et al.* 1980) and humans (Briassoulis *et al.* 2011), it is plausible to hypothesize that CORT may affect its own synthesis by a local feedback mechanism within the adrenal itself.

There is evidence that CORT can indeed affect steroid synthesis in both the adrenal and other steroidogenic organs including the gonads. With respect to the effect of CORT on the adrenal steroidogenic activity, a number of *in vitro* and *in vivo* studies have shown a decrease in adrenal responsiveness following prior exposure to stressors or ACTH. For example, experiments in cultured adrenal cells have shown that ACTH-induced CORT synthesis is rapidly inhibited (within 1–2 h) when CORT is added to the medium at high concentration (Peron *et al.* 1960, Carsia & Malamed 1979). This is in accordance with studies *in vivo* in the rat showing that previous exposure to a stressor, or to a high concentration of ACTH, inhibits CORT synthesis in response to further stimuli (Langecker & Lurie 1957, Jones & Stockham 1966), suggesting that CORT synthesis is dependent on the prior state of adrenal activity.

The molecular mechanism underlying CORT-mediated inhibition of steroidogenesis is not clear, but there is evidence that the synthetic glucocorticoid dexamethasone can inhibit the transcription of steroidogenic genes through a mechanism that involves GR (Gummow *et al.* 2006). Following ACTH stimulation and activation of CREB, StAR expression is upregulated by the CREB co-activator SF-1 and inhibited by the co-repressor DAX-1. Interestingly, ACTH can induce the transcription of SF-1 (Ragazzon *et al.* 2006) and Nur77 (Davis & Lau 1994), while inhibiting the transcription of DAX-1 (Ragazzon *et al.* 2006). Gummow *et al.* (2006) have shown that dexamethasone can induce DAX-1 transcription by a mechanism that involves the formation of an SF-1/GR complex, thus leading to inhibition of StAR transcription and ultimately to inhibition of CORT synthesis. In addition to this, CORT can repress the transcription of StAR by inhibiting both the transcription and the activity of Nur77 via a mechanism that involves active GR (Song *et al.* 2004, Martin & Tremblay 2008). In contrast to this, it has been shown that, when overexpressed *in vitro*, DAX-1 can enhance steroidogenic gene expression (Xu *et al.* 2009), and a more recent study has shown that prolonged incubation of the human adrenocortical cell line H295R with dexamethasone can indeed increase both StAR transcription and CORT production (Asser *et al.* 2014). Taken together, these studies motivate the hypothesis that dynamic secretion of CORT may be regulated by a fine balance between ACTH-mediated positive feedforward (involving upregulation and activation of steroidogenic proteins) and CORT-mediated negative feedback within the adrenal (involving GR-mediated inhibition of StAR transcriptional regulators and upregulation of StAR repressors). It is therefore

possible that pulsatile CORT synthesis observed in the adrenal in response to pulsatile ACTH stimulation may in turn be able to dynamically regulate the adrenal response to the steroidogenic stimulus. Furthermore, we have also hypothesized that when CORT secretion is elevated, as seen, for example, during a stress response or in certain conditions of disease or critical illness, this regulatory intra-adrenal feedforward–feedback mechanism may become disrupted, contributing further to abnormal CORT secretion (see the next section in this review).

To investigate this hypothesis further, in Walker *et al.* (2015) we considered the evidence for four candidate mechanisms of adrenal CORT production by studying their ability to reproduce the dynamics of the system in three experimental paradigms: i) administration of an ultradian ACTH pulse; ii) constant infusion of CRH; and iii) exposure to noise stress. Our findings provide evidence for the existence of a mechanism by which CORT can inhibit its own synthesis and/or secretion, and further that this mechanism may be transiently blocked during the early stage of an ACTH pulse. Our hypothesis is that this permits the system to respond rapidly to incoming stressors, whilst preventing uncontrolled release of glucocorticoids. The molecular basis of this ‘systems level’ mechanism remains an open question.

Because our mathematical model predicts that pulsatile CORT can modulate intra-adrenal CORT levels very rapidly, this suggests that, in addition to the genomic effects of CORT on StAR transcription already discussed, CORT can also regulate its own synthesis presumably by interfering with non-genomic mechanisms within the adrenal steroidogenic pathway, for example by regulating the activity (e.g., phosphorylation/dephosphorylation) of steroidogenic proteins including StAR and HSL (Fig. 5).

Rapid non-genomic effects of CORT at the anterior pituitary have been shown to involve annexin 1 (ANXA1), a protein that, upon CORT stimulation, is able to translocate from the cytoplasm to the outer cell surface where it exerts its regulatory effects, including inhibition of ACTH secretion from intracellular vesicles (Taylor *et al.* 1995). Interestingly, ANXA1 is also expressed in the adrenal cortex, and as observed in other tissue, its expression and activity is regulated by CORT (Davies *et al.* 2007). Remarkably, the same study also showed that ACTH stimulation of isolated adrenal gland obtained from ANXA1-null mice exhibited a greater CORT response compared to adrenal obtained from WT mice, suggesting an involvement of ANXA1 in the inhibition of CORT synthesis. Whether ANXA1 is involved in rapid CORT-mediated intra-adrenal negative




Figure 5

Model hypothesis of glucocorticoid-mediated regulation of the adrenal steroidogenic network. ACTH binding to its specific receptor MC2R induces rapid glucocorticoid secretion by PKA-mediated non-genomic regulation of steroidogenic proteins involved in cholesterol metabolism. This includes phosphorylation of hormone-sensitive lipase (HSL), a protein that increases the levels of intracellular cholesterol (the precursor of steroid hormones), and phosphorylation of steroidogenic acute regulatory protein (StAR), which promotes the transport of cholesterol into the mitochondria, where cholesterol is converted into pregnenolone by the enzyme side-chain cleavage cytochrome P450. PKA also mediates adrenal genomic activity by inducing the transcription of genes encoding for steroidogenic proteins, including StAR. Our mathematical modeling work suggests that CORT regulates an intra-adrenal inhibition mechanism of CORT synthesis, and that this is an important factor regulating CORT synthesis over the timescales of both the ultradian rhythm and the rapid response to stress. It is therefore our hypothesis that CORT may regulate the steroidogenic response by acting both at a genomic level (e.g., by regulating steroidogenic gene transcription) as well as at a non-genomic level by regulating the phosphorylation of steroidogenic proteins such as StAR and HSL.

feedback within the timescale of an ultradian pulse, however, remains to be elucidated.

In summary, our mathematical modeling work suggests that an intra-adrenal inhibition mechanism of CORT synthesis does indeed exist, and importantly, our model predictions suggest that this rapid intra-adrenal inhibition is an important factor regulating CORT synthesis over the timescales of both the basal ultradian rhythm of the HPA axis and the rapid CORT response to stress. In addition to this, our model points to the existence of a short time delay in this intra-adrenal inhibition, and that upon ACTH stimulation, this local negative feedback mechanism is rapidly antagonized, presumably via ACTH activation of the steroidogenic pathway. Our hypothesis is that these feedforward–feedback mechanisms of intra-adrenal regulation enable rapid glucocorticoid release while at the same time preventing uncontrolled release of glucocorticoids in response to large surges in ACTH associated with stress or diseased states.

Ultradian rhythm of glucocorticoids in human disease and critical illness

The release of ACTH and CORT in humans is also characterized by an ultradian rhythm with a close

temporal relationship between pulses of ACTH and CORT, as has been observed in the rat (Fig. 3D). Furthermore, consistent with changes in the pulsatile pattern observed in a number of disease models in the rat (e.g., chronic inflammation and chronic exposure to constant light), a similar disruption of cortisol pulsatility has been reported in chronic illness in humans. For example, an elevation in CORT secretion due to an increase in the mass of CORT pulses has been observed in patients affected by obstructive sleep apnea, and a complete remission of these hormonal changes was achieved following medical treatment of the condition (Henley *et al.* 2009b).

Given that the ultradian rhythm of CORT secretion is crucial for normal physiological function, it is clear that any disruption in the pulsatile pattern of hormone release could lead to impairment of CORT-regulated functions, including metabolic processes and the immune response. Indeed, maintenance of a normal ultradian rhythm of cortisol may be particularly important in major surgery and critical illness (reviewed in Gibbison *et al.* (2013)). In this regard, a recent study from our group in which the pattern of CORT secretion was investigated in patients undergoing cardiac surgery showed that a disruption in CORT secretion occurs in these patients, with high CORT levels, despite normal ACTH, observed for several hours

after the termination of the surgical procedure through the recovery period (Fig. 3A–C) (Gibbison *et al.* 2015). These data show that dissociation between ACTH and CORT occurs in these patients. It is important to highlight that in these patients both ACTH and CORT still display a pulsatile pattern of secretion, suggesting that although the adrenal sensitivity to ACTH may be altered, the mechanisms regulating pulsatility, including the dynamics of adrenal steroidogenesis, as well as the negative feedback mechanisms at the levels of the pituitary are maintained. It is also noteworthy to point out that, because a delay between the beginning of the surgery and the onset of the ACTH response was observed, it is likely that the HPA axis response in these patients may not be induced by the anesthesia or the surgical procedure itself, but other factors may be involved in the robust hormonal response observed. For example, the time of the ACTH response in these patients is consistent with the time of increase in circulating inflammatory mediators (including TNF, IL1 and IL6) that also remain elevated for about 24 h before returning to normal levels (Lahat *et al.* 1992, Roth-Isigkeit *et al.* 1999, de Mendonca-Filho *et al.* 2006). It is known that circulating cytokines can activate CORT secretion not only via inducing hypothalamic CRH and pituitary ACTH release but also by acting directly at the level of the adrenal gland (Engstrom *et al.* 2008). Indeed, cytokine receptors are expressed in the zona fasciculata of the adrenal cortex, and an increase in cytokines in the adrenal can therefore potentiate the effects of ACTH, as well as exert a direct effect on steroidogenesis, through a mechanism that involves an increase in the expression of steroidogenic genes, including StAR (Tkachenko *et al.* 2011). In addition, because pathogens can induce the expression of a number of cytokines in the adrenal gland itself through activation of toll-like receptors expressed in the adrenal, an immune-adrenal cross-talk regulating CORT synthesis has also been suggested (Judd *et al.* 2000, Zacharowski *et al.* 2006). Therefore, the mechanisms underlying the dissociation between ACTH and CORT secretion and increased adrenal sensitivity that we have observed in our clinical studies may involve circulating and/or intra-adrenal cytokines.

We further investigated the hypothesis of an involvement of inflammatory mediators in the increased adrenal sensitivity to ACTH observed in our studies using a well-established rat model of acute critical illness (administration of lipopolysaccharide (LPS)). The hormonal response to an acute injection of LPS in the rat is similar to what we observe in humans undergoing cardiac surgery. Following a rapid initial HPA axis activation, ACTH levels return to normal within 4 h of LPS administration, whereas

plasma CORT levels remain high for several hours. Since CORT secretion is tightly regulated by the pattern of ACTH under basal conditions, our data suggest that LPS injection increases adrenal sensitivity to ACTH. Interestingly, when rats are injected with a high dose of ACTH that produces plasma ACTH levels (and pattern) similar to those observed after LPS administration, CORT levels return to basal shortly after the decline of ACTH (Gibbison *et al.* 2015), further supporting our hypothesis that factors other than ACTH are responsible for the increased adrenal sensitivity observed both in our clinical and experimental studies. Consistent with the above-discussed role of cytokines in adrenal steroidogenesis, we observed that the sustained increase in CORT levels in rats injected with LPS was paralleled by increased intra-adrenal CORT levels and increased expression of steroidogenic genes including StAR and MRAP, suggesting an LPS-induced increase in adrenal steroidogenic activity. The mechanism by which cytokines affect steroidogenic gene expression has not yet been elucidated, but we found that the increase in steroidogenic gene expression was associated with a dramatic decrease in DAX-1 protein expression. In accordance with a lack of sustained increase in CORT, we did not find such effects on steroidogenic genes in rats injected with a high dose of ACTH (Spiga & Lightman 2015). In summary, both our clinical and experimental studies show that a robust dissociation between ACTH and CORT occurs in conditions of critical illness, and our data from the rat suggest that the changes in the adrenal steroidogenic pathway underlying the increase in adrenal sensitivity may not be induced by the initial peak in ACTH that occurs after surgery (human) or LPS (rat), but rather by inflammatory factors acting within the adrenal cortex to regulate CORT release.

This hypothesis is further supported by recent data from Boonen *et al.* (2014), showing that patients with sustained critical illness also have elevated levels of CORT; interestingly, the same study shows that there is a positive correlation between high levels of cytokines and high levels of CORT in these patients. Furthermore, the same investigators have observed that patients with prolonged critical illness have low levels of ACTH, presumably as a result of increased negative feedback induced by elevated cortisol levels, and this is also associated with alterations in the adrenal steroidogenic pathway including cholesterol-ester depletion as well as reduction in ACTH-regulated steroidogenic gene expression, including MC2R and StAR (Boonen *et al.* 2014). This is contrary to our hypothesis that increased adrenal responsiveness in critical illness is associated with increased steroidogenic activity in the adrenal. However, these effects on adrenal

gene expression observed in sustained critical illness are presumably the consequence of prolonged ACTH depletion observed in these patients, and are consistent with the adrenal alterations observed in mice lacking POMC, the gene that encodes for the precursor of ACTH (Karpac *et al.* 2008). Taken together, these observations suggest that in acute critical illness associated with an inflammatory response, cytokines and other immunomodulators may be responsible for elevated plasma levels of CORT. In contrast, in prolonged critical illness, high levels of plasma CORT may be due to other mechanisms, including reduced CORT metabolism (Boonen *et al.* 2013).

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

The work described in this review is funded by grants from the Medical Research Council, the Neuroendocrinology Charitable Trust and the British Heart Foundation.

References

- Apostolakis EM, Longo LD, Veldhuis JD & Yellon SM 1992 Dissociation of pulsatile cortisol and adrenocorticotropin secretion in fetal sheep during late gestation. *Endocrinology* **130** 2571–2578. (doi:10.1210/endo.130.5.1315248)
- Asser L, Hescot S, Viengchareun S, Delemer B, Trabado S & Lombes M 2014 Autocrine positive regulatory feedback of glucocorticoid secretion: glucocorticoid receptor directly impacts H295R human adrenocortical cell function. *Molecular and Cellular Endocrinology* **395** 1–9. (doi:10.1016/j.mce.2014.07.012)
- Belchetz PE, Plant TM, Nakai Y, Keogh EJ & Knobil E 1978 Hypophysial responses to continuous and intermittent delivery of hypothalamic gonadotropin-releasing hormone. *Science* **202** 631–633. (doi:10.1126/science.100883)
- Boonen E, Vervenne H, Meersseman P, Andrew R, Mortier L, Declercq PE, Vanwijngaerden YM, Spriet I, Wouters PJ, Vander Perre S *et al.* 2013 Reduced cortisol metabolism during critical illness. *New England Journal of Medicine* **368** 1477–1488. (doi:10.1056/NEJMoa1214969)
- Boonen E, Langouche L, Janssens T, Meersseman P, Vervenne H, De Samblanx E, Pironet Z, Van Dijck L, Vander Perre S, Dereese I *et al.* 2014 Impact of duration of critical illness on the adrenal glands of human intensive care patients. *Journal of Clinical Endocrinology and Metabolism* **99** 4214–4222. (doi:10.1210/jc.2014-2429)
- Briassoulis G, Damjanovic S, Xekouki P, Lefebvre H & Stratakis CA 2011 The glucocorticoid receptor and its expression in the anterior pituitary and the adrenal cortex: a source of variation in hypothalamic–pituitary–adrenal axis function; implications for pituitary and adrenal tumors. *Endocrine Practice* **17** 941–948. (doi:10.4158/EP11061.RA)
- Caraty A, Grino M, Locatelli A & Oliver C 1988 Secretion of corticotropin releasing factor (CRF) and vasopressin (AVP) into the hypophysial portal blood of conscious, unrestrained rams. *Biochemical and Biophysical Research Communications* **155** 841–849. (doi:10.1016/S0006-291X(88)80572-2)
- Carnes M, Brownfield MS, Kalin NH, Lent S & Barksdale CM 1986 Episodic secretion of ACTH in rats. *Peptides* **7** 219–223. (doi:10.1016/0196-9781(86)90216-0)
- Carnes M, Kalin NH, Lent SJ, Barksdale CM & Brownfield MS 1988a Pulsatile ACTH secretion: variation with time of day and relationship to cortisol. *Peptides* **9** 325–331. (doi:10.1016/0196-9781(88)90268-9)
- Carnes M, Lent SJ, Erisman S & Feyz J 1988b Changes in mean plasma ACTH reflect changes in amplitude and frequency of secretory pulses. *Life Sciences* **43** 1785–1790. (doi:10.1016/0024-3205(88)90277-9)
- Carnes M, Goodman BM, Lent SJ, Vo H & Jaekels R 1994 Coincident plasma ACTH and corticosterone time series: comparisons between young and old rats. *Experimental Gerontology* **29** 625–643. (doi:10.1016/0531-5565(94)90075-2)
- Caron KM, Ikeda Y, Soo SC, Stocco DM, Parker KL & Clark BJ 1997 Characterization of the promoter region of the mouse gene encoding the steroidogenic acute regulatory protein. *Molecular Endocrinology* **11** 138–147. (doi:10.1210/mend.11.2.9880)
- Carsia RV & Malamed S 1979 Acute self-suppression of corticosteroidogenesis in isolated adrenocortical cells. *Endocrinology* **105** 911–914. (doi:10.1210/endo-105-4-911)
- Churchill PF & Kimura T 1979 Topological studies of cytochromes P-450sc and P-45011 β in bovine adrenocortical inner mitochondrial membranes. Effects of controlled tryptic digestion. *Journal of Biological Chemistry* **254** 10443–10448.
- Clarke IJ & Cummins JT 1982 The temporal relationship between gonadotropin releasing hormone (GnRH) and luteinizing hormone (LH) secretion in ovariectomized ewes. *Endocrinology* **111** 1737–1739. (doi:10.1210/endo-111-5-1737)
- Dallman MF, Akana SF, Cascio CS, Darlington DN, Jacobson L & Levin N 1987a Regulation of ACTH secretion: variations on a theme of B. *Recent Progress in Hormone Research* **43** 113–173.
- Dallman MF, Akana SF, Jacobson L, Levin N, Cascio CS & Shinsako J 1987b Characterization of corticosterone feedback regulation of ACTH secretion. *Annals of the New York Academy of Sciences* **512** 402–414. (doi:10.1111/j.1749-6632.1987.tb24976.x)
- Davies E, Omer S, Buckingham JC, Morris JF & Christian HC 2007 Expression and externalization of annexin 1 in the adrenal gland: structure and function of the adrenal gland in annexin 1-null mutant mice. *Endocrinology* **148** 1030–1038. (doi:10.1210/en.2006-0732)
- Davis IJ & Lau LF 1994 Endocrine and neurogenic regulation of the orphan nuclear receptors Nur77 and Nur1 in the adrenal glands. *Molecular and Cellular Biology* **14** 3469–3483. (doi:10.1128/MCB.14.5.3469)
- Droste SK, de Groot L, Lightman SL, Reul JM & Linthorst AC 2009 The ultradian and circadian rhythms of free corticosterone in the brain are not affected by gender: an *in vivo* microdialysis study in Wistar rats. *Journal of Neuroendocrinology* **21** 132–140. (doi:10.1111/j.1365-2826.2008.01811.x)
- Engler D, Pham T, Fullerton MJ, Clarke IJ & Funder JW 1989 Evidence for an ultradian secretion of adrenocorticotropin, β -endorphin and α -melanocyte-stimulating hormone by the ovine anterior and intermediate pituitary. *Neuroendocrinology* **49** 349–360. (doi:10.1159/000125139)
- Engler D, Pham T, Liu JP, Fullerton MJ, Clarke IJ & Funder JW 1990 Studies of the regulation of the hypothalamic–pituitary–adrenal axis in sheep with hypothalamic–pituitary disconnection. II. Evidence for *in vivo* ultradian hypersecretion of proopiomelanocortin peptides by the isolated anterior and intermediate pituitary. *Endocrinology* **127** 1956–1966. (doi:10.1210/endo-127-4-1956)
- Engstrom L, Rosen K, Angel A, Fyrberg A, Mackerlova L, Konsman JP, Engblom D & Blomqvist A 2008 Systemic immune challenge activates an intrinsically regulated local inflammatory circuit in the adrenal gland. *Endocrinology* **149** 1436–1450. (doi:10.1210/en.2007-1456)
- Gibbison B, Angelini GD & Lightman SL 2013 Dynamic output and control of the hypothalamic–pituitary–adrenal axis in critical illness and major surgery. *British Journal of Anaesthesia* **111** 347–360. (doi:10.1093/bja/aet077)

- Gibbison B, Spiga F, Walker JJ, Russell GM, Stevenson K, Kershaw Y, Zhao Z, Henley D, Angelini GD & Lightman SL 2015 Dynamic pituitary–adrenal interactions in response to cardiac surgery. *Critical Care Medicine* **43** 791–800. (doi:10.1097/CCM.0000000000000773)
- Gummow BM, Scheys JO, Cancelli VR & Hammer GD 2006 Reciprocal regulation of a glucocorticoid receptor-steroidogenic factor-1 transcription complex on the Dax-1 promoter by glucocorticoids and adrenocorticotrophic hormone in the adrenal cortex. *Molecular Endocrinology* **20** 2711–2723. (doi:10.1210/me.2005-0461)
- Henley DE, Leendertz JA, Russell GM, Wood SA, Taheri S, Woltersdorf WW & Lightman SL 2009a Development of an automated blood sampling system for use in humans. *Journal of Medical Engineering & Technology* **33** 199–208. (doi:10.1080/03091900802185970)
- Henley DE, Russell GM, Douthwaite JA, Wood SA, Buchanan F, Gibson R, Woltersdorf WW, Catterall JR & Lightman SL 2009b Hypothalamic–pituitary–adrenal axis activation in obstructive sleep apnea: the effect of continuous positive airway pressure therapy. *Journal of Clinical Endocrinology and Metabolism* **94** 4234–4242. (doi:10.1210/jc.2009-1174)
- Hinz B & Hirschelmann R 2000 Rapid non-genomic feedback effects of glucocorticoids on CRF-induced ACTH secretion in rats. *Pharmaceutical Research* **17** 1273–1277. (doi:10.1023/A:1026499604848)
- Ixart G, Barbanel G, Nouguié-Soule J & Assenmacher I 1991 A quantitative study of the pulsatile parameters of CRH-41 secretion in unanesthetized free-moving rats. *Experimental Brain Research* **87** 153–158. (doi:10.1007/BF00228516)
- Ixart G, Siaud P, Mekaouche M, Barbanel G, Givalois L & Assenmacher I 1994 Short-term but not long-term adrenalectomy modulates amplitude and frequency of the CRH41 episodic release in push–pull cannulated median eminence of free-moving rats. *Brain Research* **658** 185–191. (doi:10.1016/S0006-8993(09)90025-7)
- Jasper MS & Engeland WC 1991 Synchronous ultradian rhythms in adrenocortical secretion detected by microdialysis in awake rats. *American Journal of Physiology* **261** R1257–R1268.
- Jasper MS & Engeland WC 1994 Splanchnic neural activity modulates ultradian and circadian rhythms in adrenocortical secretion in awake rats. *Neuroendocrinology* **59** 97–109. (doi:10.1159/000126645)
- Jo Y & Stocco DM 2004 Regulation of steroidogenesis and steroidogenic acute regulatory protein in R2C cells by DAX-1 (dosage-sensitive sex reversal, adrenal hypoplasia congenita, critical region on the X chromosome, gene-1). *Endocrinology* **145** 5629–5637. (doi:10.1210/en.2004-0941)
- Jones MT & Stockham MA 1966 The effect of previous stimulation of the adrenal cortex by adrenocorticotrophin on the function of the pituitary–adrenocortical axis in response to stress. *Journal of Physiology* **184** 741–750. (doi:10.1113/jphysiol.1966.sp007945)
- Jones MT, Brush FR & Neame RL 1972 Characteristics of fast feedback control of corticotrophin release by corticosteroids. *Journal of Endocrinology* **55** 489–497. (doi:10.1677/joe.0.0550489)
- Jones MT, Tiptaft EM, Brush FR, Fergusson DA & Neame RL 1974 Evidence for dual corticosteroid-receptor mechanisms in the feedback control of adrenocorticotrophin secretion. *Journal of Endocrinology* **60** 223–233. (doi:10.1677/joe.0.0600223)
- Jones MT, Hillhouse EW & Burden JL 1977 Dynamics and mechanics of corticosteroid feedback at the hypothalamus and anterior pituitary gland. *Journal of Endocrinology* **73** 405–417. (doi:10.1677/joe.0.0730405)
- Judd AM, Call GB, Barney M, McIlmoil CJ, Balls AG, Adams A & Oliveira GK 2000 Possible function of IL-6 and TNF as intraadrenal factors in the regulation of adrenal steroid secretion. *Annals of the New York Academy of Sciences* **917** 628–637. (doi:10.1111/j.1749-6632.2000.tb05428.x)
- Kalsbeek A, van der Spek R, Lei J, Ender E, Buijs RM & Fliers E 2012 Circadian rhythms in the hypothalamo–pituitary–adrenal (HPA) axis. *Molecular and Cellular Endocrinology* **349** 20–29. (doi:10.1016/j.mce.2011.06.042)
- Karpac J, Czyzewska K, Kern A, Brush RS, Anderson RE & Hochgeschwender U 2008 Failure of adrenal corticosterone production in POMC-deficient mice results from lack of integrated effects of POMC peptides on multiple factors. *American Journal of Physiology. Endocrinology and Metabolism* **295** E446–E455. (doi:10.1152/ajpendo.00762.2007)
- Kraemer FB & Shen WJ 2002 Hormone-sensitive lipase: control of intracellular tri-(di-)acylglycerol and cholesteryl ester hydrolysis. *Journal of Lipid Research* **43** 1585–1594. (doi:10.1194/jlr.R200009-JLR200)
- Lahat N, Zlotnick AY, Shtiller R, Bar I & Merin G 1992 Serum levels of IL-1, IL-6 and tumour necrosis factors in patients undergoing coronary artery bypass grafts or cholecystectomy. *Clinical and Experimental Immunology* **89** 255–260. (doi:10.1111/j.1365-2249.1992.tb06941.x)
- Langecker H & Lurie R 1957 [Inhibition of corticotropin secretion by steroids]. *Acta Endocrinologica* **25** 54–58.
- Lin D, Sugawara T, Strauss JF III, Clark BJ, Stocco DM, Saenger P, Rogol A & Miller WL 1995 Role of steroidogenic acute regulatory protein in adrenal and gonadal steroidogenesis. *Science* **267** 1828–1831. (doi:10.1126/science.7892608)
- Liu Y, Smith LI, Huang V, Poon V, Coello A, Olah M, Spiga F, Lightman SL & Aguilera G 2013 Transcriptional regulation of episodic glucocorticoid secretion. *Molecular and Cellular Endocrinology* **371** 62–70. (doi:10.1016/j.mce.2012.10.011)
- Loose DS, Do YS, Chen TL & Feldman D 1980 Demonstration of glucocorticoid receptors in the adrenal cortex: evidence for a direct dexamethasone suppressive effect on the rat adrenal gland. *Endocrinology* **107** 137–146. (doi:10.1210/endo-107-1-137)
- Mahmoud SN, Scaccianoce S, Scraggs PR, Nicholson SA, Gillham B & Jones MT 1984 Characteristics of corticosteroid inhibition of adrenocorticotrophin release from the anterior pituitary gland of the rat. *Journal of Endocrinology* **102** 33–42. (doi:10.1677/joe.0.1020033)
- Martin LJ & Tremblay JJ 2008 Glucocorticoids antagonize cAMP-induced Star transcription in Leydig cells through the orphan nuclear receptor NR4A1. *Journal of Molecular Endocrinology* **41** 165–175. (doi:10.1677/JME-07-0145)
- Martin LJ, Boucher N, Brousseau C & Tremblay JJ 2008 The orphan nuclear receptor NUR77 regulates hormone-induced Star transcription in Leydig cells through cooperation with Ca²⁺/calmodulin-dependent protein kinase I. *Molecular Endocrinology* **22** 2021–2037. (doi:10.1210/me.2007-0370)
- de Mendonça-Filho HT, Pereira KC, Fontes M, Vieira DA, de Mendonça ML, Campos LA & Castro-Faria-Neto HC 2006 Circulating inflammatory mediators and organ dysfunction after cardiovascular surgery with cardiopulmonary bypass: a prospective observational study. *Critical Care* **10** R46. (doi:10.1186/cc4857)
- Mershon JL, Sehlhorst CS, Rebar RW & Liu JH 1992 Evidence of a corticotropin-releasing hormone pulse generator in the macaque hypothalamus. *Endocrinology* **130** 2991–2996. (doi:10.1210/endo.130.5.1572307)
- Metherell LA, Chapple JP, Cooray S, David A, Becker C, Ruschendorf F, Naville D, Begeot M, Khoo B, Nurnberg P *et al.* 2005 Mutations in MRAP, encoding a new interacting partner of the ACTH receptor, cause familial glucocorticoid deficiency type 2. *Nature Genetics* **37** 166–170. (doi:10.1038/ng1501)
- Mountjoy KG, Mortrud MT, Low MJ, Simerly RB & Cone RD 1994 Localization of the melanocortin-4 receptor (MC4-R) in neuroendocrine and autonomic control circuits in the brain. *Molecular Endocrinology* **8** 1298–1308.
- Park SY, Walker JJ, Johnson NW, Zhao Z, Lightman SL & Spiga F 2013 Constant light disrupts the circadian rhythm of steroidogenic proteins in the rat adrenal gland. *Molecular and Cellular Endocrinology* **371** 114–123. (doi:10.1016/j.mce.2012.11.010)
- Peron FG, Moncloa F & Dorfman RI 1960 Studies on the possible inhibitory effect of corticosterone on corticosteroidogenesis at the adrenal level in the rat. *Endocrinology* **67** 379–388. (doi:10.1210/endo-67-3-379)
- Plotsky PM & Vale W 1985 Patterns of growth hormone-releasing factor and somatostatin secretion into the hypophysial-portal circulation of the rat. *Science* **230** 461–463. (doi:10.1126/science.2864742)

- Qian X, Droste SK, Lightman SL, Reul JM & Linthorst AC 2012 Circadian and ultradian rhythms of free glucocorticoid hormone are highly synchronized between the blood, the subcutaneous tissue, and the brain. *Endocrinology* **153** 4346–4353. (doi:10.1210/en.2012-1484)
- Ragazzon B, Lefrancois-Martinez AM, Val P, Sahut-Barnola I, Tournaire C, Chambon C, Gachancard-Bouya JL, Begue RJ, Veyssiere G & Martinez A 2006 Adrenocorticotropin-dependent changes in SF-1/DAX-1 ratio influence steroidogenic genes expression in a novel model of glucocorticoid-producing adrenocortical cell lines derived from targeted tumorigenesis. *Endocrinology* **147** 1805–1818. (doi:10.1210/en.2005-1279)
- Roth-Isigkeit A, Borstel TV, Seyfarth M & Schmucker P 1999 Perioperative serum levels of tumour-necrosis-factor alpha (TNF- α), IL-1 β , IL-6, IL-10 and soluble IL-2 receptor in patients undergoing cardiac surgery with cardiopulmonary bypass without and with correction for haemodilution. *Clinical and Experimental Immunology* **118** 242–246. (doi:10.1046/j.1365-2249.1999.01050.x)
- Rotsztein W, Lalonde J, Normand M & Fortier C 1975a Feedback inhibition of adrenocorticotropin release by corticosterone infusions in the adrenalectomized rat. *Canadian Journal of Physiology and Pharmacology* **53** 475–478. (doi:10.1139/jy75-067)
- Rotsztein WH, Normand M, Lalonde J & Fortier C 1975b Relationship between ACTH release and corticosterone binding by the receptor sites of the adenohypophysis and dorsal hippocampus following infusion of corticosterone at a constant rate in the adrenalectomized rat. *Endocrinology* **97** 223–230. (doi:10.1210/endo-97-1-223)
- Russell GM, Henley DE, Leendertz J, Douthwaite JA, Wood SA, Stevens A, Woltersdorf WW, Peeters BW, Ruijt GS, White A *et al.* 2010 Rapid glucocorticoid receptor-mediated inhibition of hypothalamic–pituitary–adrenal ultradian activity in healthy males. *Journal of Neuroscience* **30** 6106–6115. (doi:10.1523/JNEUROSCI.5332-09.2010)
- Song KH, Park YY, Park KC, Hong CY, Park JH, Shong M, Lee K & Choi HS 2004 The atypical orphan nuclear receptor DAX-1 interacts with orphan nuclear receptor Nur77 and represses its transactivation. *Molecular Endocrinology* **18** 1929–1940. (doi:10.1210/me.2004-0043)
- Spiga F & Lightman SL 2015 Dynamics of adrenal glucocorticoid steroidogenesis in health and disease. *Molecular and Cellular Endocrinology* **408** 227–234. (doi:10.1016/j.mce.2015.02.005)
- Spiga F, Harrison LR, Wood SA, Atkinson HC, MacSweeney CP, Thomson F, Craighead M, Grassie M & Lightman SL 2007 Effect of the glucocorticoid receptor antagonist Org 34850 on basal and stress-induced corticosterone secretion. *Journal of Neuroendocrinology* **19** 891–900. (doi:10.1111/j.1365-2826.2007.01605.x)
- Spiga F, Waite EJ, Liu Y, Kershaw YM, Aguilera G & Lightman SL 2011a ACTH-dependent ultradian rhythm of corticosterone secretion. *Endocrinology* **152** 1448–1457. (doi:10.1210/en.2010-1209)
- Spiga F, Liu Y, Aguilera G & Lightman SL 2011b Temporal effect of adrenocorticotrophic hormone on adrenal glucocorticoid steroidogenesis: involvement of the transducer of regulated cyclic AMP-response element-binding protein activity. *Journal of Neuroendocrinology* **23** 136–142. (doi:10.1111/j.1365-2826.2010.02096.x)
- Spiga F, Walker JJ, Terry JR & Lightman SL 2014 HPA axis-rhythms. *Comprehensive Physiology* **4** 1273–1298. (doi:10.1002/cphy.c140003)
- Stavreva DA, Wiench M, John S, Conway-Campbell BL, McKenna MA, Pooley JR, Johnson TA, Voss TC, Lightman SL & Hager GL 2009 Ultradian hormone stimulation induces glucocorticoid receptor-mediated pulses of gene transcription. *Nature Cell Biology* **11** 1093–1102. (doi:10.1038/ncb1922)
- Takemori H, Kanematsu M, Kajimura J, Hatano O, Katoh Y, Lin XZ, Min L, Yamazaki T, Doi J & Okamoto M 2007 Dephosphorylation of TORC initiates expression of the StAR gene. *Molecular and Cellular Endocrinology* **265–266** 196–204.
- Taylor AD, Loxley HD, Flower RJ & Buckingham JC 1995 Immunoneutralization of lipocortin 1 reverses the acute inhibitory effects of dexamethasone on the hypothalamo-pituitary-adrenocortical responses to cytokines in the rat *in vitro* and *in vivo*. *Neuroendocrinology* **62** 19–31. (doi:10.1159/000126984)
- Tkachenko IV, Jaaskelainen T, Jaaskelainen J, Palvimo JJ & Voutilainen R 2011 Interleukins 1 α and 1 β as regulators of steroidogenesis in human NCI-H295R adrenocortical cells. *Steroids* **76** 1103–1115. (doi:10.1016/j.steroids.2011.04.018)
- Ulrich-Lai YM & Herman JP 2009 Neural regulation of endocrine and autonomic stress responses. *Nature Reviews. Neuroscience* **10** 397–409. (doi:10.1038/nrn2647)
- Waite EJ, McKenna M, Kershaw Y, Walker JJ, Cho K, Piggins HD & Lightman SL 2012 Ultradian corticosterone secretion is maintained in the absence of circadian cues. *European Journal of Neuroscience* **36** 3142–3150. (doi:10.1111/j.1460-9568.2012.08213.x)
- Walker JJ, Terry JR & Lightman SL 2010 Origin of ultradian pulsatility in the hypothalamic–pituitary–adrenal axis. *Proceedings of the Royal Society B: Biological Sciences* **277** 1627–1633. (doi:10.1098/rspb.2009.2148)
- Walker JJ, Spiga F, Waite E, Zhao Z, Kershaw Y, Terry JR & Lightman SL 2012 The origin of glucocorticoid hormone oscillations. *PLoS Biology* **10** e1001341. (doi:10.1371/journal.pbio.1001341)
- Walker JJ, Spiga F, Gupta R, Zhao Z, Lightman SL & Terry JR 2014 Rapid intra-adrenal feedback regulation of glucocorticoid synthesis. *Journal of the Royal Society Interface* **12** article 20140875 (doi:10.1098/rsif.2014.0875)
- Watts AG, Tanimura S & Sanchez-Watts G 2004 Corticotropin-releasing hormone and arginine vasopressin gene transcription in the hypothalamic paraventricular nucleus of unstressed rats: daily rhythms and their interactions with corticosterone. *Endocrinology* **145** 529–540. (doi:10.1210/en.2003-0394)
- Widmaier EP & Dallman MF 1984 The effects of corticotropin-releasing factor on adrenocorticotropin secretion from perfused pituitaries *in vitro*: rapid inhibition by glucocorticoids. *Endocrinology* **115** 2368–2374. (doi:10.1210/endo-115-6-2368)
- Windle RJ, Wood SA, Shanks N, Lightman SL & Ingram CD 1998 Ultradian rhythm of basal corticosterone release in the female rat: dynamic interaction with the response to acute stress. *Endocrinology* **139** 443–450. (doi:10.1210/endo.139.2.5721)
- Windle RJ, Wood SA, Kershaw YM, Lightman SL, Ingram CD & Harbuz MS 2001 Increased corticosterone pulse frequency during adjuvant-induced arthritis and its relationship to alterations in stress responsiveness. *Journal of Neuroendocrinology* **13** 905–911. (doi:10.1046/j.1365-2826.2001.00715.x)
- Xu B, Yang WH, Gerin I, Hu CD, Hammer GD & Koenig RJ 2009 Dax-1 and steroid receptor RNA activator (SRA) function as transcriptional coactivators for steroidogenic factor 1 in steroidogenesis. *Molecular and Cellular Biology* **29** 1719–1734. (doi:10.1128/MCB.01010-08)
- Zacharowski K, Zacharowski PA, Koch A, Baban A, Tran N, Berkels R, Papewalis C, Schulze-Osthoff K, Knuefermann P, Zahringer U *et al.* 2006 Toll-like receptor 4 plays a crucial role in the immune-adrenal response to systemic inflammatory response syndrome. *PNAS* **103** 6392–6397. (doi:10.1073/pnas.0601527103)

Received in final form 27 May 2015

Accepted 11 June 2015

Accepted Preprint published online 11 June 2015